#### Providence

#### **Providence Digital Commons**

Providence Pharmacy PGY1 Program at Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Providence Portland and Providence St. Vincent Medical Centers 2023 Medical Centers

2023

## Standardize to Optimize: A contemplation of Pyxis, inventory, and workflow

Anthony Le Providence, anthony.le@providence.org

Scott King Providence, Scott.King2@providence.org

Follow this and additional works at: https://digitalcommons.providence.org/oaa\_ppmcstvin\_23

Fart of the Medical Education Commons, and the Pharmacy and Pharmaceutical Sciences Commons

#### **Recommended Citation**

Le, Anthony and King, Scott, "Standardize to Optimize: A contemplation of Pyxis, inventory, and workflow" (2023). *Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Medical Centers 2023.* 10.

https://digitalcommons.providence.org/oaa\_ppmcstvin\_23/10

This is brought to you for free and open access by the Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Medical Centers at Providence Digital Commons. It has been accepted for inclusion in Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Medical Centers 2023 by an authorized administrator of Providence Digital Commons. For more information, please contact digitalcommons@providence.org.



# Standardize to Optimize: A Contemplation of Pyxis, Inventory, and Workflow

Anthony Le, PharmD; Scott King, PharmD, BSN



## Background

 Automated dispensing cabinets (ADC) are commonly used within hospitals to store and distribute medications in a timely manner.

•ADC use creates a more efficient workflow and decreases medication errors through unit dose drawers and pockets within ADCs.

•Optimization of ADCs has yet to be fully defined. The paucity in data has created a knowledge gap on standard optimization methods.

		Change in Vend to Fi	ll Ratio	
	Change in vend-to-fill ratio, by medication			
	•			
200 -	•			
	1			StationName
				Station Name: CARD-A8
100 -		•		Station Name: CARD-B8
		•		Station Name: CARD-B8

	• •		
DISCL	JSSION		

<u>Results</u>

#### Change in Vend to Fill Ratio

 Inversed relationship with vend to fill ratio with increasing PAR

 Primarily increase in number of refills with experimental group

### Change in Number of Vends

Decreased among all ADCs

•ADC medication stockouts can create delay in medication administration, increase pharmacy refill times, and decrease pharmacy productivity.

 Additional costs and reduced storage space are associated with unused medications in automated dispensing cabinets.

#### Purpose

Evaluate ADC efficiency through periodic automatic replacement (PAR) level adjustments.
Correlate number of vends and number of fills with increased medication minimum and maximum levels.

## Objectives

#### Primary outcomes

• Vend to fill ratios of the top 50 medications per Pyxis machine.



 Non-significant change between control versus experimental group

#### Change in Number of Refills

Ratio of 1.88 increase in refills by increasing PAR minimums one additional day
Strong inversed relationship suggests minimum PAR levels are triggered more frequently in top 50 used medications.

## **Optimization Data Limitations**

•Rx Auditor data was extrapolated beyond 90 days.
•Current medication usage may fluctuate within different ADCs.

- Only top 50 medications were evaluated and correlations cannot be extrapolated to less frequently used medications.
- Limited data on medication stockouts and expired medications

# Going Forward

Secondary outcomes
Number of medication refills
Number of medication vends

## Methods

•Study design

Prospective study

Inclusion criteria

 Top 50 medications used from each Pyxis from two different hospital services (3 Pyxis machines per service).

Identified through RxAuditor 90-day report.
Service lines with similar ADC usage

Exclusion criteria

- Medications with lower number of vends outside of the top 50 medications for each service
- Medications with multiple pockets in the same ADC
  Non-unit dosed medications
- Interventions

 Removal of unused, non-essential medications from ADC



Number of Refills

#### Change in number of refills, by medication



Continue with current minimum medication PAR levels
Validate complete ADC medication standardization
Combine medications with multiple pockets into one standard pocket for each ADC
Validate correlation with reduced number of PAR levels to test efficiency

Complete analysis of all medication usage loaded into ADCs
Complete analysis of medications loaded into ADCs from various services throughout the hospital
Complete a detailed statistical analysis on medication stockouts, variations, medication returns in relations to PAR maximum level adjustment

## References

 Tony Vu, Yifan She; Automated Dispensing Cabinet Optimization in a Level 2 Trauma Center. *Journal of Contemporary Pharmacy Practice* 1 September 2020; 67 (4): 17–21. doi: https://doi.org/10.37901/jcphp20-00005
 O'Neil DP, Miller A, Cronin D, Hatfield CJ. A comparison of automated dispensing cabinet optimization methods. Am J Health Syst Pharm. 2016

## Increase PAR minimum levels for one hospital service's top 50 ADC medications

# Jul 1;73(13):975-80. doi: 10.2146/ajhp150423. Epub 2016 May 23. PMID: 27217517.

3. Mann J, Dipper A, Rowe E, Hatfield C. Lean six sigma in the pharmacy department. Pharmacy Times. Sept 2014; 3(5).