Providence

Providence Digital Commons

2021 Providence Nurse Research Conference

Providence Nursing Research Conference

6-2021

Exploring the Relationship Between COVID-19 Unit Designation and Nurse Burnout Syndrome

Gisele Bazan Covenant Medical Center, BazanGN@covhs.org

Tiffany Patterson Covenant Children's Hospital, tpatterson@covhs.org

Kelsey Sawyer Covenant Medical Center, jollyk1@covhs.org

Jamie Roney Covenant Medical Center, jroney@covhs.org

Erin Whitley

See next page for additional authors

Follow this and additional works at: https://digitalcommons.providence.org/prov_rn_conf_all

Part of the Pediatric Nursing Commons

Recommended Citation

Bazan, Gisele; Patterson, Tiffany; Sawyer, Kelsey; Roney, Jamie; Whitley, Erin; Mihandoust, Sahar; and Long, JoAnn, "Exploring the Relationship Between COVID-19 Unit Designation and Nurse Burnout Syndrome" (2021). *2021 Providence Nurse Research Conference*. 17. https://digitalcommons.providence.org/prov_rn_conf_all/17

This is brought to you for free and open access by the Providence Nursing Research Conference at Providence Digital Commons. It has been accepted for inclusion in 2021 Providence Nurse Research Conference by an authorized administrator of Providence Digital Commons. For more information, please contact digitalcommons@providence.org.

Authors

Gisele Bazan, Tiffany Patterson, Kelsey Sawyer, Jamie Roney, Erin Whitley, Sahar Mihandoust, and JoAnn Long

This is available at Providence Digital Commons: https://digitalcommons.providence.org/prov_rn_conf_all/17

Exploring the Relationship Between COVID-19 Unit Designation and Nurse Burnout Syndrome

Gisele Bazan, BSN, RN, CCRN-K Tiffany Patterson, BSN, RN Kelsey Sawyer, BSN, RN, CCRN-K Jamie Roney, DNP, RN-BC, CCRN-K Erin Whitley, MSN, RN-BC Sahar Mihandoust, PhD, M.Arch. JoAnn Long, PhD, RN, NEA-BC



Learning Objective

Learners will be able to describe the impact of COVID-19 pandemic care to nurse burnout in a tertiary care center in the southwest United States.





Introduction

The COVID-19 pandemic impacted healthcare due to surges in infected patient and respiratory failure.

(Bateman et al., 2020; González-Gil et al., 2020; Lasater et al., 2020; Morgantini et al., 2020; Xinghuang Liu et al., 2020)

Nursing burnout syndrome (NBS) results from occupational factors leading to mental health problems.

(Giusti et al., 2020; Hu et al., 2020; Kok et al., 2019; Meltzer et al, 2004; Mihandoust et al., 2021; Morgantini et al., 2020; Moss et al., 2016; Murat et al., 2020; Poncet et al., 2007; Salazar et al., 2020; Sanghera et al., 2020; Sayilan et al., 2020; Xinghuang Liu et al., 2020)



Background

Intensive care units (ICUs) dedicated to caring for COVID-19 infected individuals may be more susceptible to NBS due to:

- -increased burdens of futile care
- -high mortality rates
- overstretched healthcare systems

(Giusti et al., 2020; Hu et al., 2020; Kok et al., 2019; Meltzer et al, 2004; Mihandoust et al., 2021; Morgantini et al., 2020; Moss et al., 2016; Murat et al., 2020; Poncet et al., 2007; Salazar et al., 2020; Sanghera et al., 2020; Sayilan et al., 2020; Xinghuang Liu et al., 2020)



Purpose

The purpose of this study was to compare burnout levels in ICU nurses providing direct care to COVID infected patients with nurses caring for non-COVID infected patients.



Significance



Identifying sociodemographic, workrelated, and psychological predictors of NBS may help organizations mitigate, or at least minimize, the negative psychological impact on ICU nurses working during future pandemics.



Methods

A comparative descriptive study was conducted:

- 1. Neonatal and pediatric ICU to cardiac and medical ICU survey scores measuring predictors of NBS
- 2. NBS scores for nurses working in six critical care units captured in 2019 prior to the COVID-19 pandemic to scores captured in a ICU after conversion to dedicated COVID-19 ICUs in 2020



Methods

Maslach Burnout Inventory (MBI) used

Subscales:

- Emotional Exhaustion (EE)
- Depersonalization (DP)
- Personal
 Accomplishment (PA)





Maslach Burnout Inventory Scales

Emotional Exhaustion (EE)

• 9-item scale measures feelings of being emotionally overextended and exhausted by one's work.

Depersonalization (DP)

• 5-item scale measures an unfeeling and impersonal response toward recipients of one's service, care, treatment, or instruction.

Personal Accomplishment (PA)

• 8-item scale measures feelings of competence and successful achievement in one's work.





Methods

A convenience sample of direct care registered nurses working in four ICUs in a large tertiary care hospital participated in the study.

Female nurses working day shifts were surveyed in February and March 2019 in six critical care units.

In November 2020, four ICUs were used for sampling of both male and female nurses working either day or night 12-hour shifts based on designation as COVID-19 ICUs.



Maslach Burnout Inventory Scales



Data Analysis

- Data entered into SPSS version 27
- Descriptive, frequencies, & Q-Q plots conducted Assumptions of normality met in EE and DP subscales
 - PA subscale data was skewed in both comparison groups
- Independent sample t-tests compared differences in mean scores on EE and DP subscales between COVID-19 and Non-COVID-19 units
- Mann-Whitney U test compared differences in mean PA scores between independent groups

CovenantHea

Demographics





Demographics

Total years in nursing

< 2 years = 22% (20)

2-5 years = 29% (27)

6-10 years = 15% (14)

11-15 years = 8% (7)

16-20 years = 4% (4)

21-25 years = 8% (7)

26-30 years = 2% (2) >30 years = 10% (9)

Highest level of education

28%(26) Diploma

10%(9) ADN

57%(52) BSN

3%(3) MSN



Extra Shifts Worked





COVID UNITS – AVERAGE 1.17 (SD = 1.09) EXTRA SHIFTS WORKED

NON-COVID UNITS – AVERAGE 0.28 (SD = 0.65) EXTRA SHIFTS WORKED



Results

We hypothesized burnout scores among nurses working in non-COVID ICUs would be less than burnout scores for nurses working in designated COVID ICUs.

	COVID- designated units (n=52)	Non-COVID units (n=38)	p-value
EE	18.2 (moderate)	8.6 (low)	0.0001
DP	12.8 (high)	5.0 (low)	0.0001
PA	32.6 (high)	36.8 (moderate)	0.08





Independent Sample Test – COVID and Non-COVID ICU's

	Levene's Test for Equality of Variance			t-test for Equality of Means			95% Confidence Interval of Diff		
Sub- scale	F	Sig.	t	Df	Sig. (2-tailed)	Mean Diff	Std. Error Mean	Lower	Upper
EE (Equal variance assumed)	8.55	.004	5.06	88	.0001	9.63	1.90	5.85	13.44
EE (Equal variance not assumed)			5.31	87.99	.0001	9.63	1.81	6.03	13.24
DP (Equal variance assumed)	31.52	.0001	4.82	88	.0001	8.03	1.66	4.72	11.35
DP (Equal variance not assumed)			5.31	78.51	.0001	8.03	1.51	5.02	11.05



Results

Independent Sample ttest results support a statistically significant difference in EE and DP scores between COVID and Non-COVID designated units.





Large effect size for both EE and DP scores supports important differences in the magnitude and strength of scores between groups.

Cohen's d	Standardizer	Point	95% Confidence Interval		
		Estimate	Lower	Upper	
EE	8.91	1.08	.630	1.52	
DP	7.80	1.03	.582	1.47	

• Nurses working on COVID-designated units experienced more EE and DP than nurses working on non-COVID units.





The Mann Whitney U test was used to examine the difference between the non-normally distributed mean PA scores in the COVID and non-COVID units.

COVID compared to Non-COVID scores	Mann- Whitney U Test Statistic	Standard Error	Standardized Test Statistic	Asymptotic Sig. (2 sided)
PA	1198.00	122.29	1.71	0.08

• A non-statistically significant difference between mean PA scores in the COVID compared to Non-COVID designated units was noted.





Results

We hypothesized nurse burnout scores in six critical care (CC) units measured in 2019 (Pre-COVID) would be less than burnout scores in 2020 in the primary COVID-19 ICU.

	Pre-COVID critical care RN's (N=51)	COVID critical care RN's (N=23)	p-value
EE	15.41 (low)	18.2 (moderate)	.59
DP	10.29 (high)	12.8 (high)	.74

- Statistical significance was not noted pre and post COVID-19 pandemic using independent sample t-test
- Clinical significance was noted by an increase (indicating more burnout)
- EE sub-scale (increased from low pre to moderate)

Independent Sample Test – Pre-COVID CC Units & COVID ICU

	Levene's Test for Equality of Variance			t-test for Equality of Means			95% Confidence Interval of Diff		
Sub- scale	F	Sig.	t	Df	Sig. (2-tailed)	Mean Diff	Std. Error Mean	Lower	Upper
EE (Equal variance assumed)	.392	.533	.537	72	.593	1.28	2.38	-3.47	6.04
EE (Equal variance not assumed)			5.23	39.84	.604	1.28	2.45	-3.68	6.24
DP (Equal variance assumed)	1.08	.301	.331	72	.742	.61	1.87	-3.11	4.35
DP (Equal variance not assumed)			.305	35.53	.762	.61	2.02	-3.49	4.73



Independent Samples Pre-COVID CC Units & COVID ICU Effect Size

Effect size for both EE and DP scores did not support differences in the magnitude and strength of scores between Pre-COVID and COVID designated units.

Cohen's d	Standardizer	Point	95% Confidence Interval	
		Estimate	Lower	Upper
EE	9.51	.135	358	.627
DP	7.45	.083	410	.575



Discussion

Important differences in **EE and DP scores** between COVID and non-**COVID** units were supported. Further research is required to establish relationships between sociodemographic and workrelated psychological predictors of NBS.

Understanding relationships between variables may guide development of strategies to build nurse resilience and decrease NBS in ICU settings impacted during future pandemics.



Conclusion



NBS has been identified as a global problem facing ICU clinicians.



Pinpointing associations between COVID-19 infection and nurse burnout may lead to strategies to mitigate burnout in those caring for individuals during future pandemics.



References

- Bateman, M. E., Hammer, R., Byrne, A., Ravindran, N., Chiurco, J., Lasky, S., Denson, R., Brown, M., Myers, L., Zu, Y., & Denson, J. L. (2020). Death Cafés for prevention of burnout in intensive care unit employees: study protocol for a randomized controlled trial (STOPTHEBURN). *Trials*, 21(1), 1–9. https://doi.org/10.1186/s13063-020-04929-4
- Hu, D., Kong, Y., Li, W., Han, Q., Zhang, X., Zhu, L. X., Wan, S. W., Liu, Z., Shen, Q., Yang, J., He, H.-G., & Zhu, J. (2020). Frontline nurses' burnout, anxiety, depression, and fear statuses and their associated factors during the COVID-19 outbreak in Wuhan, China: A large-scale cross-sectional study. *EClinicalMedicine*, 24. https://doi.org/10.1016/j.eclinm.2020.100424
- Kok N, van Gurp J, Teerenstra S, van der Hoeven H, Fuchs M, Hoedemaekers C, Zegers M. Coronavirus Disease 2019 Immediately Increases Burnout Symptoms in ICU Professionals: A Longitudinal Cohort Study. *Crit Care Med*. 2021 Jan 28. doi: 10.1097/CCM.00000000004865. Epub ahead of print. PMID: 33555778.
- Lasater KB, Aiken LH, Sloane DM, French R, Martin B, Reneau K, Alexander M, McHugh MD. Chronic hospital nurse understaffing meets COVID-19: an observational study. *BMJ Qual Saf*. 2020 Aug 18:bmjqs-2020-011512. doi: 10.1136/bmjqs-2020-011512. Epub ahead of print. PMID: 32817399; PMCID: PMC7443196.
- Meltzer LS, & Huckabay LM. (2004). Critical care nurses' perceptions of futile care and its effect on burnout. American Journal of Critical Care, 13(3), 202–208. https://doi.org/10.4037/ajcc2004.13.3.202
- Mihandoust, S., Pati, D., Lee, J., & Roney, J. K. (2021). Exploring the relationship between visual access to nature and nurse burnout. [Manuscript accepted for publication]. *Health Environments Research & Design Journal.*

Morgantini LA, Naha U, Wang H, Francavilla S, Acar Ö, Flores JM, Crivellaro S, Moreira D, Abern M, Eklund M, Vigneswaran HT, Weine SM. Factors Contributing to Healthcare Professional Burnout During the COVID-19 Pandemic: A Rapid Turnaround Global Survey. medRxiv [Preprint]. 2020 May 22:2020.05.17.20101915. doi: 10.1101/2020.05.17.20101915. Update in: PLoS One. 2020 Sep 3;15(9):e0238217. PMID: 32511501; PMCID: PMC7273269.



References

- Moss, M., & Good, V. S. (2016). An Official Critical Care Societies Collaborative Statement: Burnout Syndrome in Critical Care Health Care Professionals: A Call for Action. *American Journal of Critical Care*, 25(4), 368–376. https://doi.org/10.4037/ajcc2016133
- Murat M, Köse S, Savaşer S. Determination of stress, depression and burnout levels of front-line nurses during the COVID-19 pandemic. *Int J Ment Health Nurs*. 2020 Nov 21:10.1111/inm.12818. doi: 10.1111/inm.12818. Epub ahead of print. PMID: 33222350; PMCID: PMC7753629.
- Poncet MC, Toullic P, Papazian L, Kentish-Barnes N, Timsit J, Pochard F, Chevret S, Schlemmer B, & Azoulay E. (2007). Burnout syndrome in critical care nursing staff. *American Journal of Respiratory & Critical Care Medicine*, 175(7), 698–704. https://doi.org/10.1164/rccm.200606-806oc
- Suleiman, M. N., Gomez, U. J. L., Aguayo, E. R., Cañadas, D. L. F. G. A., De La Fuente, S. E. I., & Albendín, G. L. (2020). The effect of mindfulness training on burnout syndrome in nursing: A systematic review and meta-analysis. *Journal of Advanced Nursing* (John Wiley & Sons, Inc.), 76(5), 1124–1140. https://doi.org/10.1111/jan.14318
- Xinghuang Liu, Jie Chen, Dongke Wang, Xin Li, Erchuan Wang, Yu Jin, Yanling Ma, Cheng Yu, Chang Luo, Lei Zhang, Chuang Liu, Yangshiyu Zhou, Ling Yang, Jun Song, Tao Bai, & Xiaohua Hou. (2020). COVID-19 Outbreak Can Change the Job Burnout in Health Care Professionals. *Frontiers in Psychiatry*, *11*. https://doi.org/10.3389/fpsyt.2020.563781

