

Providence St. Joseph Health
Providence St. Joseph Health Digital Commons

Books, Presentations, Posters, Etc.

6-2019

Ibuprofen Does Not Increase Blood Pressure in Preeclampsia

Sofia Costas

Providence St. Vincent Medical Center, Portland, Oregon, sofia.costas@providence.org

Sherry Hutton

Providence St. Vincent Medical Center, Portland, Oregon, sherry.hutton@providence.org

Cindy Kenyon

Providence St. Vincent Medical Center, Portland, Oregon, cindy.kenyon@providence.org

Follow this and additional works at: https://digitalcommons.psjhealth.org/other_pubs



Part of the [Maternal, Child Health and Neonatal Nursing Commons](#)

Recommended Citation

Costas, Sofia; Hutton, Sherry; and Kenyon, Cindy, "Ibuprofen Does Not Increase Blood Pressure in Preeclampsia" (2019). *Books, Presentations, Posters, Etc.*. 88.

https://digitalcommons.psjhealth.org/other_pubs/88

This Poster is brought to you for free and open access by Providence St. Joseph Health Digital Commons. It has been accepted for inclusion in Books, Presentations, Posters, Etc. by an authorized administrator of Providence St. Joseph Health Digital Commons. For more information, please contact digitalcommons@providence.org.

Ibuprofen Does Not Increase Blood Pressure in Preeclampsia

Sofia Costas BSN, RNC, Sherry Hutton BSN, RN, Cindy Kenyon BSN, RNC
 Providence St. Vincent Medical Center, Portland, Oregon



Objective

The aim of this IRB approved study was to evaluate the impact of ibuprofen on mean arterial pressure (MAP) in the immediate postpartum period of women with preeclampsia.

Background

In 2013, The American College of Obstetricians and Gynecologists (ACOG) made a recommendation to withhold ibuprofen in postpartum preeclamptic women.

While the United States is facing an opioid epidemic, preeclamptic postpartum women are receiving a greater amount of narcotics for pain control because ibuprofen is not considered safe to use in the postpartum period. The belief is that ibuprofen increases blood pressure in postpartum women with preeclampsia.

Contact Information:
 Sofia Costas Sofia.Costas@providence.org
 Sherry Hutton Sherry.Hutton@providence.org
 Cindy Kenyon Cindy.Kenyon@providence.org

Methods

A query identified 633 preeclamptic perinatal patients who delivered at Providence St. Vincent Medical Center, from January 2017 to December 2017, who had severe hypertension (HTN) (BP $\geq 160/105$) during their hospital stay.

A total of 169 patients met our criteria, 66 (39%) received ibuprofen and 103 (61%) did not. A stratified random sample selected 60 preeclamptic postpartum patients, including 30 who received ibuprofen postpartum and 30 who did not.


Inclusion Criteria:

- Preeclampsia diagnosis
- Magnesium sulfate during the postpartum period
- Treatment for severe HTN after delivery

Exclusion Criteria:

- Patients presenting with a BP $\geq 160/105$ at 2 hours following delivery


The baseline MAP was obtained at 2 hours postpartum and the MAPs for each 24 hour interval were averaged. The MAPs were compared in women who received ibuprofen versus those who did not. T-tests were used to determine statistical significance between the MAPs of the two groups of patients at the following hourly intervals: 24, 48, 72 and 96. Power calculations were done to ensure adequate sample size.



Graph 1.

Results

The average MAP scores are displayed in Figure 1 and Graph 1. When patients who did receive ibuprofen were compared with patients who did not receive ibuprofen, there was no statistical significant differences in MAP comparisons (baseline [p=0.46], 2 to 24 hours [p=0.46], 24 to 48 hours [p=0.56], 48 to 72 hours [p=0.45], 72 to 96 hours [p=0.49], average of total hours [p=0.48]).



Graph 2.

Implications for Nursing Practice

Based on our findings, ibuprofen following birth is safe for patients with postpartum preeclampsia and should be included in the pain control plan in order to reduce narcotic use.

Our results demonstrated that there was no significant difference in MAP for preeclampsia patients who received ibuprofen in the postpartum period compared to the patients who did not receive ibuprofen. This finding is consistent with current literature.

Since the completion of our study, the Perinatal Special Care Unit at Providence St. Vincent Medical Center has increased the percentage of ibuprofen administered to preeclamptic postpartum women from 39% to 95%. Graph 2.

Key limitations of this study included the limited sample size that met criteria. Demographic data was not collected on the patients included in this study.

Average MAP Values with Standard Deviation

Ibuprofen	2 Hour Baseline	2 to 24 Hours	24-<48 Hours	48-<72 Hours	72-96 Hours	Total Average
NO	98.6 \pm 7.1	97.1 \pm 6.4	95.1 \pm 7.4	98.0 \pm 8	97.8 \pm 6.4	97.2 \pm 7.3
YES	96.8 \pm 9.9	95.9 \pm 6.4	95.6 \pm 7.4	96.4 \pm 7.0	97.7 \pm 6.8	96.5 \pm 7.5
p	0.46	0.46	0.56	0.45	0.49	0.48

Figure 1.

References

- Blue, N. R., Drake-Lavelle, S et al. (2018). LB04: Effect of Ibuprofen versus acetaminophen on postpartum hypertension in preeclampsia with severe features: A double-masked, randomized controlled trial. *American Journal of Obstetrics and Gynecology*, 218(1). doi:10.1016/j.ajog.2017.11.599
- Gracia, P. V., Solis, V., & Ortega, N. (2016). Ibuprofen versus acetaminophen as a post-partum analgesic for women with severe pre-eclampsia: Randomized clinical study. *The Journal of Maternal-Fetal & Neonatal Medicine*, 30(11), 1279-1282. doi:10.1080/14767058.2016.1210599
- Hirshberg, J. S., Fatima, S., López, J. D., Tuuli, M. G., Macones, G. A., & Cahill, A. G. (2018). 335: Preeclampsia with severe features: Is postpartum ibuprofen use associated with the need for antihypertensive medication. *American Journal of Obstetrics and Gynecology*, 218(1). doi:10.1016/j.ajog.2017.10.271
- Viteri, O. A., England, J. A et al. (2017). Association of Nonsteroidal Antiinflammatory Drugs and Postpartum Hypertension in Women With Preeclampsia With Severe Features. *Obstetrics & Gynecology*, 130(4), 830-835. doi:10.1097/aog.0000000000002247
- Wasden, S. W., Ragsdale, E. S., Chasen, S. T., & Skupski, D. W. (2014). Impact of non-steroidal anti-inflammatory drugs on hypertensive disorders of pregnancy. *Pregnancy Hypertension: An International Journal of Womens Cardiovascular Health*, 4(4), 259-263. doi:10.1016/j.preghy.2014.06.001